



STIC Search Report

EIC 1700

STIC Database Tracking Number: 182904

**TO: Eisa Elhilo
Location: REM 9A60
Art Unit : 1751
March 22, 2006**

Case Serial Number: 10/798454

**From: Les Henderson
Location: EIC 1700
REM 4B28 / 4A30
Phone: 571-272-2538**

Leslie.henderson@uspto.gov

Search Notes



STIC Search Results Feedback Form

EIC17000

Questions about the scope or the results of the search? Contact *the EIC searcher* or contact:

Kathleen Fuller, EIC 1700 Team Leader
571/272-2505 REMSEN 4B28

Voluntary Results Feedback Form

- I am an examiner in Workgroup: Example: 1713
➤ Relevant prior art **found**, search results used as follows:

- ☐ 102 rejection
- ☐ 103 rejection
- ☐ Cited as being of interest.
- ☐ Helped examiner better understand the invention.
- ☐ Helped examiner better understand the state of the art in their technology.

Types of relevant prior art found:

- ☐ Foreign Patent(s)
- ☐ Non-Patent Literature
(journal articles, conference proceedings, new product announcements etc.)

➤ Relevant prior art **not found**:

- ☐ Results verified the lack of relevant prior art (helped determine patentability).
- ☐ Results were not useful in determining patentability or understanding the invention.

Comments:

Drop off or send completed forms to EIC1700 REMSEN 4B28

=> d his

(FILE 'HOME' ENTERED AT 13:19:22 ON 22 MAR 2006)

FILE 'HCAPLUS' ENTERED AT 13:20:34 ON 22 MAR 2006

L1 E US20050081313/PN
1 S US20050081313/PN
SEL RN

FILE 'REGISTRY' ENTERED AT 13:21:54 ON 22 MAR 2006

L2 53 S E1-E53

FILE 'LREGISTRY' ENTERED AT 13:25:58 ON 22 MAR 2006

L3 STR

FILE 'REGISTRY' ENTERED AT 13:36:05 ON 22 MAR 2006

L4 0 S L3
L5 SCR 2043
L7 0 S L3 NOT L5 FUL

FILE 'LREGISTRY' ENTERED AT 13:38:25 ON 22 MAR 2006

L8 STR L3
L9 STR L8

FILE 'REGISTRY' ENTERED AT 14:01:17 ON 22 MAR 2006

L10 0 S L9
L11 0 S L9 NOT L5

FILE 'LREGISTRY' ENTERED AT 14:03:21 ON 22 MAR 2006

L12 STR L9

FILE 'REGISTRY' ENTERED AT 14:05:10 ON 22 MAR 2006

L13 1 S L12
L14 102 S L12 FUL
SAV L14 LEE454/A
L15 39 S L14 AND L2

FILE 'LREGISTRY' ENTERED AT 14:11:23 ON 22 MAR 2006

L16 STR L12

FILE 'REGISTRY' ENTERED AT 14:12:36 ON 22 MAR 2006

L17 1 S L16 SSS SAM SUB=L14
L18 85 S L16 SSS FUL SUB=L14
SAV L18 LEE454A/A
L19 STR L16

FILE 'REGISTRY' ENTERED AT 14:15:33 ON 22 MAR 2006


L20 1 S L19 SSS SAM SUB=L14
L21 53 S L19 SSS FUL SUB=L14
SAV L21 LEE454B/A
L22 24 S L21 AND L2
L23 39 S L18 AND L2

FILE 'HCAPLUS' ENTERED AT 14:17:49 ON 22 MAR 2006

L24 7 S L23
L25 2 S L22
L26 13 S L21
L27 24 S L18
L28 28 S L14
L29 7 S L24 OR L15
L30 1 S L1 AND L26
L31 15 S L28 NOT L26

=> d que stat 126


L12 STR



VAR G2=C/N/O
 REP G3=(0-2) C
 NODE ATTRIBUTES:
 DEFAULT MLEVEL IS ATOM
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
 RING(S) ARE ISOLATED OR EMBEDDED
 NUMBER OF NODES IS 14

STEREO ATTRIBUTES: NONE
 L14 102 SEA FILE=REGISTRY SSS FUL L12
 L19 STR



VAR G1=NH2/NO2
 VAR G2=C/N/O
 REP G3=(0-2) C
 VAR G4=X/AK
 NODE ATTRIBUTES:
 DEFAULT MLEVEL IS ATOM
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
 RING(S) ARE ISOLATED OR EMBEDDED
 NUMBER OF NODES IS 15

STEREO ATTRIBUTES: NONE
 L21 53 SEA FILE=REGISTRY SUB=L14 SSS FUL L19
 L26 13 SEA FILE=HCAPLUS ABB=ON PLU=ON L21

=> => d 126 1-13 ibib abs hitstr hitind

L26 ANSWER 1 OF 13 HCAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 2004:758628 HCAPLUS
 DOCUMENT NUMBER: 141:282402
 TITLE: Hair dye composition comprising a heterocyclic
 oxidation base and a 2,3,5-triaminopyridine

INVENTOR(S): coupler
 Kravtchenko, Sylvain; Lagrange, Alain; Vidal,
 Laurent; Fadli, Aziz
 PATENT ASSIGNEE(S): L'Oreal, Fr.
 SOURCE: Fr. Demande, 29 pp.
 CODEN: FRXXBL
 DOCUMENT TYPE: Patent
 LANGUAGE: French
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 2852241	A1	20040917	FR 2003-3115	2003 0313
EP 1459732	A1	20040922	EP 2004-290634	2004 0309

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,
 MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ,
 EE, HU, PL, SK, HR

PRIORITY APPLN. INFO.: FR 2003-3115 A
 2003
 0313

OTHER SOURCE(S): MARPAT 141:282402


AB The title compns. are used for the dyeing of keratinous fibers. A hair dye composition contained 7-dimethylamino-3-aminopyrazolopyrimidine 2x10⁻³, 3,5-diamino-2-pyrrolidinopyridine 2x10⁻³, benzyl alc. 2, ethoxylated polyethylene glycol 3, ethanol 18, Oramix CG110 10, sodium metabisulfite 0.205, sequestering agent q.s., and water q.s. 100 g.

IT 756498-17-6, N-(3,5-Diaminopyridin-2-yl)-2-methylpyrrolidine 756498-41-6 756498-45-0
 756498-48-3 756498-50-7 756498-56-3
 756498-59-6

RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
 (hair dye composition comprising heterocyclic oxidation base and 2,3,5-triaminopyridine coupler)


RN 756498-17-6 HCAPLUS

CN 3,5-Pyridinediamine, 2-(2-methyl-1-pyrrolidinyl)- (9CI) (CA INDEX NAME)



RN 756498-41-6 HCAPLUS


CN 3,5-Pyridinediamine, 2-(2,5-dimethyl-1-pyrrolidinyl)- (9CI) (CA INDEX NAME)



RN 756498-45-0 HCAPLUS


CN L-Proline, 1-(3,5-diamino-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 756498-48-3 HCAPLUS

CN 2-Pyrrolidinecarboxamide, 1-(3,5-diamino-2-pyridinyl)- (9CI) (CA INDEX NAME)




RN 756498-50-7 HCAPLUS

CN 2-Pyrrolidinecarboxamide, 1-(3,5-diamino-2-pyridinyl)-N,N-dimethyl- (9CI) (CA INDEX NAME)




RN 756498-56-3 HCAPLUS

CN 3,5-Pyridinediamine, 2-[3-(dimethylamino)-1-pyrrolidinyl]- (9CI) (CA INDEX NAME)



RN 756498-59-6 HCAPLUS
 CN 3,5-Pyridinediamine, 2-(4-methyl-1-piperazinyl)- (9CI) (CA INDEX NAME)



IC ICM A61K007-13
 CC 62-3 (Essential Oils and Cosmetics)
 IT 51-35-4, 4-Hydroxyproline 109-01-3 110-85-0, Piperazine, biological studies 110-89-4, Piperidine, biological studies 111-49-9, Homopiperidine 123-75-1, Pyrrolidine, biological studies 147-85-3, Proline, biological studies 498-63-5, 2-Hydroxymethylpyrrolidine 504-03-0, 2,6-Dimethylpiperidine 535-75-1, 2-Carboxypiperidine 567-36-2, 3-Hydroxyproline 765-38-8, 2-Methylpyrrolidine 2812-47-7, 2-Pyrrolidinecarboxamide 3378-71-0, 2,5-Dimethylpyrrolidine 3433-37-2, 2-Hydroxymethylpiperidine 4318-37-0, N-Methyl homopiperazine 4606-65-9, 3-Hydroxymethylpiperidine 5227-53-2 5382-16-1, 4-Hydroxypiperidine 5626-66-4, 2,5-Pyrrolidinedimethanol 6859-99-0, 3-Hydroxypiperidine 19889-77-1, 2-Piperidinecarboxamide 27230-48-4, 3-Hydroxy-2-hydroxymethylpiperidine 40499-83-0, 3-Hydroxypyrrolidine 53427-65-9 67523-79-9D, derivs. 69478-75-7, 3-Dimethylaminopyrrolidine 79286-79-6, 3-Aminopyrrolidine 83030-08-4, 3-Methylaminopyrrolidine 89364-91-0, 3-Hydroxy-2-carboxamidopyrrolidine 99319-03-6, 2,4-Pyrrolidinedicarboxylic acid 128508-51-0 130497-29-9 169750-98-5 178105-25-4 188925-57-7 433980-61-1 473541-96-7, 3,4-Dihydroxypyrrolidine 756498-17-6, N-(3,5-Diaminopyridin-2-yl)-2-methylpyrrolidine 756498-39-2, N-(3,5-Diaminopyridin-2-yl)pyrrolidine 756498-41-6 756498-45-0 756498-46-1 756498-48-3 756498-50-7 756498-52-9 756498-54-1 756498-56-3 756498-58-5 756498-59-6 757967-87-6 757967-88-7 757967-89-8 757967-90-1 757967-91-2
 RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
 (hair dye composition comprising heterocyclic oxidation base and 2,3,5-triaminopyridine coupler)

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L26 ANSWER 2 OF 13 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2004:753132 HCAPLUS

DOCUMENT NUMBER: 141:265564

TITLE: Coupling agents having a 2,3,5-triaminopyridine structure and their use for dyeing keratinic fibers

INVENTOR(S): Fadli, Aziz; Vidal, Laurent

PATENT ASSIGNEE(S): L'oreal, Fr.

SOURCE: Eur. Pat. Appl., 36 pp.

CODEN: EPXXDW

DOCUMENT TYPE: Patent

LANGUAGE: French

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1457199	A1	20040915	EP 2004-290611	2004 0305
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR				
FR 2852240	A1	20040917	FR 2003-3114	2003 0313
JP 2004277423	A2	20041007	JP 2004-71608	2004 0312
US 2005081313	A1	20050421	US 2004-798454	2004 0312
PRIORITY APPLN. INFO.:				
			FR 2003-3114	A 2003 0313
			US 2003-467124P	P 2003 0502

OTHER SOURCE(S): MARPAT 141:265564.


AB Hair dye compns. comprising 2,3,5-triaminopyridine structure are used as coupling agent for dyeing hair. Thus, 2-(3,5-dimethylpiperidin-1-yl)pyridine-3,5-diamine (I) was prepared by the reduction of 2-(3,5-dimethylpiperidin-1-yl)-3,5-dinitropyridine on palladium/charcoal. A hair dye composition contained I 10-3 mole, 2-[(4-Amino-phenyl)-(2-hydroxyethyl)-amino]-ethanol sulfate 10-3 mole, and water and excipients q.s. 100 g.

IT 756498-17-6 756498-41-6 756498-43-8
756498-45-0 756498-48-3 756498-50-7
756498-56-3 756498-59-6

RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(coupling agents having triaminopyridine structure and their use for dyeing keratinic fibers)


RN 756498-17-6 HCAPLUS

CN 3,5-Pyridinediamine, 2-(2-methyl-1-pyrrolidinyl)- (9CI) (CA INDEX NAME)




RN 756498-41-6 HCAPLUS

CN 3,5-Pyridinediamine, 2-(2,5-dimethyl-1-pyrrolidinyl)- (9CI) (CA INDEX NAME)




RN 756498-43-8 HCAPLUS
 CN 2-Pyrrolidinemethanol, 1-(3,5-diamino-2-pyridinyl)- (9CI) (CA INDEX NAME)




RN 756498-45-0 HCAPLUS
 CN L-Proline, 1-(3,5-diamino-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.




RN 756498-48-3 HCAPLUS
 CN 2-Pyrrolidinecarboxamide, 1-(3,5-diamino-2-pyridinyl)- (9CI) (CA INDEX NAME)




RN 756498-50-7 HCAPLUS
 CN 2-Pyrrolidinecarboxamide, 1-(3,5-diamino-2-pyridinyl)-N,N-dimethyl- (9CI) (CA INDEX NAME)



RN 756498-56-3 HCAPLUS
 CN 3,5-Pyridinediamine, 2-[3-(dimethylamino)-1-pyrrolidinyl]- (9CI)
 (CA INDEX NAME)




RN 756498-59-6 HCAPLUS
 CN 3,5-Pyridinediamine, 2-(4-methyl-1-piperazinyl)- (9CI) (CA INDEX NAME)



IT 756498-18-7P 756498-20-1P 756498-23-4P
 756498-25-6P 756498-27-8P 756498-29-0P
 756498-31-4P 756498-33-6P
 RL: COS (Cosmetic use); SPN (Synthetic preparation); BIOL
 (Biological study); PREP (Preparation); USES (Uses)
 (coupling agents having triaminopyridine structure and their
 use for dyeing keratinic fibers)
 RN 756498-18-7 HCAPLUS
 CN Methanol, compd. with 2-(2-methyl-1-pyrrolidinyl)-3,5-
 pyridinediamine hydrochloride (5:5:9) (9CI) (CA INDEX NAME)

CM 1

CRN 756498-17-6
 CMF C10 H16 N4




CM 2

CRN 67-56-1
 CMF C H4 O


H₃C-OH

RN 756498-20-1 HCAPLUS
 CN 3,5-Pyridinediamine, 2-(2,5-dimethyl-1-pyrrolidinyl)-,
 dihydrochloride (9CI) (CA INDEX NAME)




● 2 HCl

RN 756498-23-4 HCAPLUS
 CN 3,5-Pyridinediamine, 2-(4-methyl-1-piperazinyl)-, dihydrochloride
 (9CI) (CA INDEX NAME)




● 2 HCl

RN 756498-25-6 HCAPLUS
 CN 3,5-Pyridinediamine, 2-(4-ethyl-1-piperazinyl)-, trihydrochloride
 (9CI) (CA INDEX NAME)




● 3 HCl

RN 756498-27-8 HCAPLUS
 CN 3,5-Pyridinediamine, 2-(3,5-dimethyl-1-piperidinyl)-,
 dihydrochloride (9CI) (CA INDEX NAME)




●2 HCl

RN 756498-29-0 HCAPLUS
 CN 2-Piperidineethanol, 1-(3,5-diamino-2-pyridinyl)-, dihydrochloride
 (9CI) (CA INDEX NAME)




●2 HCl

RN 756498-31-4 HCAPLUS
 CN 4-Piperidinecarboxamide, 1-(3,5-diamino-2-pyridinyl)-,
 dihydrochloride (9CI) (CA INDEX NAME)



●2 HCl

RN 756498-33-6 HCAPLUS
 CN 3,5-Pyridinediamine, 2-(2-methyl-1-piperidinyl)-, dihydrochloride
 (9CI) (CA INDEX NAME)



● 2 HCl

IT 756498-19-8P 756498-21-2P 756498-24-5P

756498-26-7P 756498-28-9P 756498-30-3P

756498-32-5P 756498-34-7P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP

(Preparation); RACT (Reactant or reagent)

(coupling agents having triaminopyridine structure and their use for dyeing keratinic fibers)


RN 756498-19-8 HCAPLUS

CN Pyridine, 2-(2-methyl-1-pyrrolidinyl)-3,5-dinitro- (9CI) (CA INDEX NAME)




RN 756498-21-2 HCAPLUS

CN Pyridine, 2-(2,5-dimethyl-1-pyrrolidinyl)-3,5-dinitro- (9CI) (CA INDEX NAME)




RN 756498-24-5 HCAPLUS

CN Piperazine, 1-(3,5-dinitro-2-pyridinyl)-4-methyl- (9CI) (CA INDEX NAME)




RN 756498-26-7 HCAPLUS


CN Piperazine, 1-(3,5-dinitro-2-pyridinyl)-4-ethyl- (9CI) (CA INDEX NAME)




RN 756498-28-9 HCAPLUS
 CN Pyridine, 2-(3,5-dimethyl-1-piperidinyl)-3,5-dinitro- (9CI) (CA INDEX NAME)




RN 756498-30-3 HCAPLUS
 CN 2-Piperidineethanol, 1-(3,5-dinitro-2-pyridinyl)- (9CI) (CA INDEX NAME)



RN 756498-32-5 HCAPLUS
 CN 4-Piperidinecarboxamide, 1-(3,5-dinitro-2-pyridinyl)- (9CI) (CA INDEX NAME)



RN 756498-34-7 HCAPLUS
 CN Pyridine, 2-(2-methyl-1-piperidinyl)-3,5-dinitro- (9CI) (CA INDEX NAME)



IC ICM A61K007-13
ICS C07D401-04; C07D413-04
CC 62-3 (Essential Oils and Cosmetics)
IT 756498-17-6 756498-39-2 756498-41-6
756498-43-8 756498-45-0 756498-46-1
756498-48-3 756498-50-7 756498-52-9
756498-54-1 756498-56-3 756498-58-5
756498-59-6
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(coupling agents having triaminopyridine structure and their
use for dyeing keratinic fibers)
IT 756498-14-3P 756498-15-4P 756498-18-7P
756498-20-1P 756498-22-3P 756498-23-4P
756498-25-6P 756498-27-8P 756498-29-0P
756498-31-4P 756498-33-6P 756498-36-9P
756498-38-1P
RL: COS (Cosmetic use); SPN (Synthetic preparation); BIOL
(Biological study); PREP (Preparation); USES (Uses)
(coupling agents having triaminopyridine structure and their
use for dyeing keratinic fibers)
IT 90871-11-7P 134787-63-6P 573987-10-7P 756498-16-5P
756498-19-8P 756498-21-2P 756498-24-5P
756498-26-7P 756498-28-9P 756498-30-3P
756498-32-5P 756498-34-7P 756498-37-0P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP
(Preparation); RACT (Reactant or reagent)
(coupling agents having triaminopyridine structure and their
use for dyeing keratinic fibers)
REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE
FOR THIS RECORD. ALL CITATIONS AVAILABLE
IN THE RE FORMAT

L26 ANSWER 3 OF 13 HCAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 2003:354713 HCAPLUS
DOCUMENT NUMBER: 139:164474
TITLE: Kinetic and equilibrium studies of
 σ -adduct formation and nucleophilic
substitution in the reactions of
2-phenoxy-3,5-dinitropyridine and
2-ethoxy-3,5-dinitropyridine with aliphatic
amines in dipolar aprotic solvents
AUTHOR(S): Crampton, Michael R.; Emokpae, Thomas A.;
Howard, Judith A. K.; Isanbor, Chukwuemeka;
Mondal, Raju
CORPORATE SOURCE: Chemistry Department, Durham University,
Durham, DH1 3LE, UK
SOURCE: Organic & Biomolecular Chemistry (2003), 1(6),
1004-1011
CODEN: OBCRAK; ISSN: 1477-0520
PUBLISHER: Royal Society of Chemistry
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 139:164474
AB The reactions of aliphatic amines with 2-phenoxy-3,5-dinitropyridine,
4, and 2-ethoxy-3,5-dinitropyridine, 5, in DMSO result in the
rapid reversible formation of anionic σ -adducts at the

6-position. Kinetic studies show that proton transfer from the initially formed zwitterions to base may be rate-limiting. Slower reactions result, except in the case of 5 and piperidine, in displacement of the 2-substituent via intermediates which have lower thermodyn. stabilities than their 6-isomers. Base catalysis of the substitution process is attributed in the case of 4 to rate-limiting proton transfer from zwitterionic intermediates, but in 5 to acid catalysis of ethoxide departure (SB-GA mechanism). X-ray crystallog. of 5 shows a planar nonstrained structure although the structure of 2-piperidino-3,5-dinitropyridine, 10c, shows distortion resulting from steric interactions of the 2- and 3-substituents. Kinetic and equilibrium results are compared with those for related reactions of the more sterically strained 2,4,6-trinitrobenzene derivs. Results for the reactions of 4 and 5 with pyrrolidine in three dipolar aprotic solvents are compared. Values of equilibrium consts. for σ -adduct formation decrease in the order DMSO > DMF >> acetonitrile, while values of rate consts. for proton transfer are in the reverse order.


IT 573987-11-8 573987-12-9

RL: CPS (Chemical process); FMU (Formation, unclassified); PEP (Physical, engineering or chemical process); PRP (Properties); RCT (Reactant); FORM (Formation, nonpreparative); PROC (Process); RACT (Reactant or reagent)

(kinetics and equilibrium of σ -adduct formation and nucleophilic substitution in reactions of 2-phenoxy- and 2-ethoxy-3,5-dinitropyridine with aliphatic amines in dipolar aprotic solvents)


RN 573987-11-8 HCAPLUS

CN Pyridine, 6-ethoxy-1,2-dihydro-3,5-dinitro-2-(1-pyrrolidinyl)-, ion(1-) (9CI) (CA INDEX NAME)



RN 573987-12-9 HCAPLUS

CN Pyridine, 6-ethoxy-1,2-dihydro-3,5-dinitro-2-(1-piperidinyl)-, ion(1-) (9CI) (CA INDEX NAME)



CC 22-12 (Physical Organic Chemistry)

Section cross-reference(s): 75

IT 573987-11-8 573987-12-9

RL: CPS (Chemical process); FMU (Formation, unclassified); PEP (Physical, engineering or chemical process); PRP (Properties); RCT (Reactant); FORM (Formation, nonpreparative); PROC (Process); RACT (Reactant or reagent)

(kinetics and equilibrium of σ -adduct formation and nucleophilic substitution in reactions of 2-phenoxy- and

2-ethoxy-3,5-dinitropyridine with aliphatic amines in dipolar aprotic solvents)

REFERENCE COUNT: 37 THERE ARE 37 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L26 ANSWER 4 OF 13 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1986:608919 HCAPLUS

DOCUMENT NUMBER: 105:208919

TITLE: Quinazoline derivatives and antihypertensive preparations containing them

INVENTOR(S): Yokoyama, Keiichi; Kato, Koji; Kitahara, Takumi; Ohno, Hiroyasu; Nishina, Takashi; Awaya, Akira; Nakano, Takuo; Watanabe, Kazuyuki; Saruta, Sakae; Kumakura, Mikio

PATENT ASSIGNEE(S): Mitsui Petrochemical Industries, Ltd., Japan; Mitsui Pharmaceuticals, Inc.

SOURCE: Eur. Pat. Appl., 235 pp.

CODEN: EPXXDW

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 188094	A2	19860723	EP 1985-309049	1985 1212
EP 188094	A3	19871223		
EP 188094	B1	19920318		
R: DE, FR, GB, IT				
JP 61140568	A2	19860627	JP 1984-263015	1984 1214
JP 05028709	B4	19930427		
JP 62056488	A2	19870312	JP 1985-194968	1985 0905
JP 03071430	B4	19911113		
JP 62067077	A2	19870326	JP 1985-204463	1985 0918
JP 05029223	B4	19930428		
PRIORITY APPLN. INFO.:			JP 1984-263015	A 1984 1214
			JP 1985-194968	A 1985 0905
			JP 1985-204463	A 1985 0918

OTHER SOURCE(S): CASREACT 105:208919; MARPAT 105:208919
GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT
*

AB Piperazinyl- and homopiperazinylquinazolines I (R1 = H, MeO; R2, R3 = H, alkoxy; R4 = H, NH2; R5 = substituted 2-pyrimidinyl, 2-pyridinyl, 2-quinolinyl, fused pyrimidinyl; n = 2, 3) were prepared as antihypertensives. Thus, 4-benzyl-1-piperazinecarboxamide sulfate was cyclocondensed with MeCOC(CO2Me):CHOME to give pyrimidinecarboxylate II. This was amidated with EtNH2 and cyclocondensed with DMF to give pyridopyrimidinone III, which was debenzylated and condensed with 4-amino-2-chloro-6,7-dimethoxyquinazoline to give piperazinylquinazoline IV. In rats 1 mg IV/kg orally reduced blood pressure 23.0% after 6 h, the effect lasting 24 h. Tablets were prepared each containing I 1, starch 60, microcrystn. cellulose 35, light silica 3, and Mg stearate 1 mg.


IT 104965-86-8P 104987-66-8P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of, as antihypertensive)


RN 104965-86-8 HCAPLUS

CN 4-Quinazolinamine, 2-[4-(3,5-dinitro-2-pyridinyl)-1-piperazinyl]-6,7-dimethoxy- (9CI) (CA INDEX NAME)



RN 104987-66-8 HCAPLUS

CN 4-Quinazolinamine, 2-[4-(3,5-dinitro-2-pyridinyl)-1-piperazinyl]-6,7-dimethoxy-, hydrochloride (9CI) (CA INDEX NAME)



●x HCl

IC ICM C07D471-04

ICS C07D487-04; C07D239-84; C07D495-04; C07D401-12; A61K031-495; A61K031-505

ICI C07D471-04, C07D239-00, C07D221-00; C07D487-04, C07D239-00, C07D209-00; C07D487-04, C07D243-00, C07D239-00; C07D495-04, C07D333-00, C07D239-00; C07D487-04

CC 28-16 (Heterocyclic Compounds (More Than One Hetero Atom))
Section cross-reference(s): 1, 63

IT	102839-00-9P	104964-10-5P	104964-11-6P	104964-12-7P
	104964-13-8P	104964-14-9P	104964-15-0P	104964-16-1P
	104964-17-2P	104964-18-3P	104964-19-4P	104964-20-7P
	104964-21-8P	104964-22-9P	104964-23-0P	104964-24-1P
	104964-25-2P	104964-26-3P	104964-27-4P	104964-28-5P
	104964-29-6P	104964-30-9P	104964-31-0P	104964-32-1P
	104964-33-2P	104964-34-3P	104964-35-4P	104964-36-5P
	104964-37-6P	104964-38-7P	104964-39-8P	104964-40-1P
	104964-41-2P	104964-42-3P	104964-43-4P	104964-44-5P

104964-45-6P	104964-46-7P	104964-47-8P	104964-48-9P
104964-49-0P	104964-50-3P	104964-51-4P	104964-52-5P
104964-53-6P	104964-54-7P	104964-55-8P	104964-56-9P
104964-57-0P	104964-58-1P	104964-59-2P	104964-60-5P
104964-61-6P	104964-62-7P	104964-63-8P	104964-64-9P
104964-65-0P	104964-66-1P	104964-67-2P	104964-68-3P
104964-69-4P	104964-70-7P	104964-71-8P	104964-72-9P
104964-73-0P	104964-74-1P	104964-75-2P	104964-76-3P
104964-77-4P	104964-78-5P	104964-79-6P	104964-80-9P
104964-81-0P	104964-82-1P	104964-83-2P	104964-84-3P
104964-85-4P	104964-86-5P	104964-87-6P	104964-88-7P
104964-89-8P	104964-90-1P	104964-91-2P	104964-92-3P
104964-93-4P	104964-94-5P	104964-95-6P	104964-96-7P
104964-97-8P	104964-98-9P	104964-99-0P	104965-00-6P
104965-01-7P	104965-02-8P	104965-03-9P	104965-04-0P
104965-05-1P	104965-06-2P	104965-07-3P	104965-08-4P
104965-09-5P	104965-10-8P	104965-11-9P	104965-12-0P
104965-13-1P	104965-14-2P	104965-15-3P	104965-16-4P
104965-17-5P	104965-18-6P	104965-19-7P	104965-20-0P
104965-21-1P	104965-22-2P	104965-23-3P	104965-24-4P
104965-25-5P	104965-26-6P	104965-27-7P	104965-28-8P
104965-29-9P	104965-30-2P	104965-31-3P	104965-32-4P
104965-33-5P	104965-34-6P	104965-35-7P	104965-36-8P
104965-37-9P	104965-38-0P	104965-39-1P	104965-40-4P
104965-41-5P	104965-42-6P	104965-43-7P	104965-44-8P
104965-45-9P	104965-46-0P	104965-47-1P	104965-48-2P
104965-49-3P	104965-50-6P	104965-51-7P	104965-52-8P
104965-53-9P	104965-54-0P	104965-55-1P	104965-56-2P
104965-57-3P	104965-58-4P	104965-59-5P	104965-60-8P
104965-61-9P	104965-62-0P	104965-63-1P	104965-64-2P
104965-65-3P	104965-66-4P	104965-67-5P	104965-68-6P
104965-69-7P	104965-70-0P	104965-71-1P	104965-72-2P
104965-73-3P	104965-74-4P	104965-75-5P	104965-76-6P
104965-77-7P	104965-78-8P	104965-79-9P	104965-80-2P
104965-81-3P	104965-82-4P	104965-83-5P	104965-84-6P
104965-85-7P	104965-86-8P	104965-87-9P	104965-88-0P
104965-89-1P	104965-90-4P	104965-91-5P	104965-92-6P
104987-58-8P	104987-59-9P	104987-60-2P	104987-61-3P
104987-62-4P	104987-63-5P	104987-64-6P	104987-65-7P
104987-66-8P	105010-30-8P	105201-43-2P	105201-44-3P

RL: BAC (Biological activity or effector, except adverse); BSU
(Biological study, unclassified); SPN (Synthetic preparation); THU
(Therapeutic use); BIOL (Biological study); PREP (Preparation);
USES (Uses)
(preparation of, as antihypertensive)

L26 ANSWER 5 OF 13 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1983:488003 HCAPLUS

DOCUMENT NUMBER: 99:88003

TITLE: Nucleophilic reaction upon electron-deficient
pyridone derivatives. V. Anionic
 σ -adducts of 1-methyl-3,5-dinitro-2-
pyridone

AUTHOR(S): Tohda, Yasuo; Ariga, Masahiro; Kawashima,
Toshihide; Matsumura, Eizo


CORPORATE SOURCE: Dep. Chem., Osaka Kyoiku Univ., Osaka, 543,
Japan

SOURCE: Chemistry Letters (1983), (5), 715-18
CODEN: CMLTAG; ISSN: 0366-7022

DOCUMENT TYPE: Journal

LANGUAGE: English

GI



AB Alkylammonium salts I of anionic σ -adducts of 1-methyl-3,5-dinitro-2-pyridone with amine and C nucleophiles were isolated and characterized. The C adducts are more stable than the amine adducts, which dissociate in dilute MeOH. The acetone adduct was converted into 4-nitro-1-pyrrolidinylbenzene quant. by heating with pyrrolidine.

IT 86670-27-1P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)


RN 86670-27-1 HCAPLUS

CN 2(1H)-Pyridinone, 5,6-dihydro-1-methyl-3,5-dinitro-6-(1-pyrrolidinyl)-, compd. with pyrrolidine (1:2) (9CI) (CA INDEX NAME)

CM 1

CRN 86670-26-0

CMF C10 H14 N4 O5



CM 2

CRN 123-75-1

CMF C4 H9 N



CC 27-16 (Heterocyclic Compounds (One Hetero Atom))

IT 10220-22-1P 86653-20-5P 86653-22-7P 86653-24-9P

86653-26-1P 86653-32-9P 86653-35-2P 86653-37-4P

86653-39-6P 86670-27-1P 86714-58-1P 86714-59-2P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)

L26 ANSWER 6 OF 13 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1979:203958 HCAPLUS

DOCUMENT NUMBER: 90:203958

TITLE: Products of the reaction of benzimidazole derivatives with 2-chloro-3,5-dinitropyridine


AUTHOR(S): Zakhs, E. R.; Subbotina, M. A.; El'tsov, A. V.

CORPORATE SOURCE: USSR

SOURCE: Zhurnal Organicheskoi Khimii (1979), 15(1),

DOCUMENT TYPE:
LANGUAGE:
OTHER SOURCE(S):
GI

200-6
CODEN: ZORKAE; ISSN: 0514-7492
Journal
Russian
CASREACT 90:203958




AB Benzimidazoles I (R1 = 3,5-dinitro-2-pyridyl, Me; R2 = H, Me; R3 = CN, H, CO2Me) were prepared in 50-90% yields by treating benzimidazoleacetonitriles with 2-chloro-3,5-dinitropyridine, follows by methylation, hydration, esterification, or protonation (to give the endocyclic onium compds.).

IT 70309-18-1P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation and isomerism of)

RN 70309-18-1 HCAPLUS


CN 2-Pyridineacetonitrile, α -[1-(3,5-dinitro-2-pyridinyl)-1,3-dihydro-2H-benzimidazol-2-ylidene]-3,5-dinitro- (9CI) (CA INDEX NAME)



IT 70309-19-2P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and protonation of)

RN 70309-19-2 HCAPLUS

CN Acetonitrile, [1-(3,5-dinitro-2-pyridinyl)-1,3-dihydro-2H-benzimidazol-2-ylidene]- (9CI) (CA INDEX NAME)



CC 28-10 (Heterocyclic Compounds (More Than One Hetero Atom))

IT 70309-18-1P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation and isomerism of)

IT 70309-19-2P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP
(Preparation); RACT (Reactant or reagent)
(preparation and protonation of)

L26 ANSWER 7 OF 13 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1970:78819 HCAPLUS

DOCUMENT NUMBER: 72:78819

TITLE: Synthesis and properties of
2-fluoro-3,5-dinitro-4-methyl-pyridine and
2-fluoro-3,5-dinitro-6-methylpyridine

AUTHOR(S): Talik, Tadeusz; Talik, Zofia

CORPORATE SOURCE: Wyzsza Szk. Ekon., Wroclaw, Pol.

SOURCE: Roczniki Chemii (1969), 43(11), 1961-70


CODEN: ROCHAC; ISSN: 0035-7677

DOCUMENT TYPE: Journal

LANGUAGE: German

AB The title compds. were synthesized via the following products:
2-nitramino-5-nitro-4-picoline (I), 2-amino-3,5-dinitro-4-picoline
(II), 2-fluoro-3,5-dinitro-4-picoline (III); and
2-nitramino-5-nitro-6-picoline (IV), 2-amino-3,5-dinitro-6-
picoline (V) and 2-fluoro-3,5-dinitro-6-picoline (VI). The
fluorine atom in III and VI was easily substituted in reactions
with H⁺-H₂O, alcs., amines, and amino acids. Thus, 10 g I or IV
(optionally containing some of the 3-NO₂ isomer) in 50 ml concentrated H₂SO₄
was heated 1 hr on a steam bath to give 7.1 g II, m. 183°,
or 7.7 g V, m. 179°, (both from EtOH-Me₂NCHO), resp. II or
V (6.0 g) was dissolved in a solution of HBF₄ (from 30 g H₃BO₃ and 75
ml 38% HF aqueous), 9 g NaNO₂ slowly added at 0°, and the mixture
kept 20 min at room temperature to yield 2.7 g III, m. 33°, or
2.5 g VI, b₃ 122-3°, m. 35°, resp. III or VI was
refluxed in 5 ml 1:1 HCl-H₂O 10 min to give 95.2%
2-hydroxy-3,5-dinitro-4-picoline, m. 254° (decomposition)
(H₂O-Me₂NCHO), or 98.5% 2-hydroxy-3,5-dinitro-6-picoline m.
238° (decomposition), (H₂O-EtOH), resp. III or VI (0.5 g) was
refluxed with 5 ml MeOH or EtOH 90min to give 94.6%
2-methoxy-3,5-dinitro-4-picoline, m. 129°; 95.5%
2-ethoxy-3,5-dinitro-4-picoline, m. 87°; 94.4%
2-methoxy-3,5-dinitro-6-picoline, m. 54°; and
2-ethoxy-3,5-dinitro-6-picoline, m. 63°, resp. (all
recrystd. from aqueous EtOH). To 0.5 g II in 3 ml EtOH was added 3 ml
saturated NH₃, or gaseous amine solution in EtOH, or 3 ml liquid amine to
give R- in 2-(R-substituted)-3,5-dinitro-4-picolines (VII) (R, %
yield, and m.p. given): NH₂, 98.5, 187° (H₂O-Me₂-HCHO);
NH₂Et, 99.2, 103° (H₂O-EtOH); NHCH₂CH₂OH, 95.9, 118°
(H₂O-EtOH); NHPH, 88.2, 156° (C₆H₆-EtOH); NHNHPH, 95.4,
126° (decomposition) (EtOH); NMe₂, 88.9, 74° (EtOH); and
NEt₂, 94.9, 87° (H₂O-EtOH). From VI by the above procedure
were obtained 2-(R-substituted)-3,5-dinitro-6-picolines (VIII)
(same data given): NH₂, 99.1, 179° (EtOH-H₂O); NH₂Et, 89.3,
80° (EtOH-H₂O); NHCH₂CH₂OH, 98.4, 105° (EtOH-H₂O);
NHPH, 95.1, 163° (C₆H₆-EtOH); NHNHPH, 95.5, 150°
(EtOH-Me₂NCHO); NMe₂, 95.4, 84° (EtOH); and NEt₂, 82.1,
52° (EtOH-H₂O). III (0.005 mole) in 10 ml EtOH was added
to a solution of an amino acid (0.005 mole in 5 ml H₂O and 0.01 mole
NaHCO₃), and the mixture kept 10 min at room temperature to give the
following VII (same data given): NHCHMeCO₂H-DL, 80.2, 166°
(H₂O-EtOH); NHCH(CH₂OH)CO₂H-DL, 77.5, 160° (decomposition)
(H₂O-EtOH); NHCH(CH₂CH₂SMe)CO₂H-L, 92.4, 163° (decomposition)
(EtOH-Me₂NCHO); and NHCH(CH₂CO₂H)CO₂H-L, 90.5, 162°
(decomposition) (H₂O-EtOH). Similarly VI gave the following VIII (same
data given): NHCH₂CO₂H, 95.5, 209° (decomposition)
(H₂O-Me₂NCHO); NHCHMeCO₂H-DL, 82.1, 121° (EtOH-H₂O);
NHCH(CH₂CONH₂)CO₂H-L, 78.5, 180° (H₂O-EtOH); and

IT N-(L-hydroxyprolinyl), 78.5, 55° (H₂O-EtOH).
 25864-44-2P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of)
 RN 25864-44-2 HCAPLUS
 CN Proline, 5-hydroxy-1-(6-methyl-3,5-dinitro-2-pyridyl)- (8CI) (CA
 INDEX NAME)




CC 27 (Heterocyclic Compounds (One Hetero Atom))
 IT 25782-38-1P 25782-39-2P 25782-40-5P 25782-41-6P
 25782-42-7P 25782-43-8P 25782-44-9P 25782-45-0P
 25782-46-1P 25864-29-3P 25864-30-6P 25864-31-7P
 25864-32-8P 25864-33-9P 25864-34-0P 25864-35-1P
 25864-36-2P 25864-37-3P 25864-38-4P 25864-39-5P
 25864-40-8P 25864-41-9P 25864-42-0P 25864-43-1P
 25864-44-2P 25979-16-2P 25979-17-3P 26169-30-2P
 30505-22-7P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of)

L26 ANSWER 8 OF 13 HCAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1969:47456 HCAPLUS
 DOCUMENT NUMBER: 70:47456
 TITLE: 4-Nitroimidazoles
 INVENTOR(S): Klink, Rainer; Hepding, Ludwig
 PATENT ASSIGNEE(S): Merck, E., A.-G.
 SOURCE: Brit., 5 pp.
 CODEN: BRXXAA
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
GB 1133408		19681113	GB 1967-38544	1967 0822
DE 1620043			DE	
DE 1695397			DE	
FR 7260			FR	
US 3491105		19700120	US	1967 0925
ZA 6905397		19690000	ZA	
PRIORITY APPLN. INFO.:			DE	1966 1015
			DE	1967 0301

OTHER SOURCE(S): MARPAT 70:47456

- GI For diagram(s), see printed CA Issue.
- AB The title compds. of formula I with antitrichomonadal activity, are prepared by alkylation of 2-substituted-4(5)-nitroimidazoles. Thus, a mixture of 2.3 g. Na in 100 ml. EtOH, 11.3 g. 4(5)-nitroimidazole, and 17 g. 2-chloro-5-nitropyridine was refluxed 15 min. and poured into H₂O to give 19 g. I (R₁ = H, R₂ = 5-nitro-2-pyridyl), m. 196-8° (Me₂CO-petroleum ether). Heating a mixture of 15 g. 2-methyl-4(5)-nitroimidazole Na salt (II), 15.7 g. p-O₂NC₆H₄Cl, and 100 ml. ProH in a sealed tube 30 min. at 150° gave 18.5 g. I (R₁ = Me, R₂ = p-O₂NC₆H₄) (III). Similarly, refluxing a mixture of 15 g. II, 14.1 g. p-O₂NC₆H₄F (IV), and 100 ml. Me₂NCHO 15 min. gave 15 g. III, m. 185-7° (Me₂CO). The following I were similarly prepared (R₁, R₂, and m.p. given): H, 3-nitro-2-pyridyl, 144-6°; H, 3,5-dinitro-2-pyridyl, 146-9°; H, p-O₂NC₆H₄, 188-90°; H, o-O₂NC₆H₄, 123-5°; Me, o-O₂NC₆H₄, 177-9°; H, 2,4-(O₂N)₂C₆H₃, 154-6°; Me, 2,4-(O₂N)₂C₆H₃, 194-6°; H, 2,4-Me-(O₂N)₂C₆H₃, 185-7°; Me, 2,4-Me(O₂N)₂C₆H₃, 196-7°; Me, 3,4-Me(O₂N)₂C₆H₃, 176-8°; Me, 5-nitro-2-pyridyl, 175-8°; Me, 3-nitro-2-pyridyl, 179-82°; and Me, 3,5-dinitro-2-pyridyl, 150-2°. Heating a mixture of 101.5 g. 1-(4-nitrophenyl)-2-methylimidazole (V) (m. 134-5°) (obtained by stirring 2-methylimidazole and IV 2 hrs. at 120-5° in Me₂NCHO), 200 ml. 65% aqueous HNO₃, and 50 ml. concentrated H₂SO₄ 1 hr. at 120° with portionwise addition of 150 ml. concentrated H₂SO₄ and working up gave 5.1 g. III and 16 g. 1-(4-nitrophenyl)-2-methyl-5-nitroimidazole, m. 164.5-5.5°; 19.5 g. V was recovered.
- IT 21722-03-2P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)
- RN 21722-03-2 HCAPLUS
- CN Pyridine, 2-(2-methyl-4-nitroimidazol-1-yl)-3,5-dinitro- (8CI)
(CA INDEX NAME)



- IC C07D
- CC 28 (Heterocyclic Compounds (More Than One Hetero Atom))
- IT 21721-89-1P 21721-90-4P 21721-91-5P 21721-92-6P
21721-93-7P 21721-94-8P 21721-95-9P 21721-96-0P
21721-97-1P 21721-98-2P 21721-99-3P 21722-00-9P
21722-01-0P 21722-02-1P 21722-03-2P 21722-04-3P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)

L26 ANSWER 9 OF 13 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1965:480971 HCAPLUS

DOCUMENT NUMBER: 63:80971

ORIGINAL REFERENCE NO.: 63:14976f-g

TITLE: The intramolecular participation of the pyridyl group in the acid hydrolysis of dinitropyridyl dipeptides

AUTHOR(S): Signor, Angelo; Bordignon, Emilio

CORPORATE SOURCE: Univ. Padua, Italy

SOURCE: Journal of Organic Chemistry (1965), 30(10), 3447-51

CODEN: JOCEAH; ISSN: 0022-3263

DOCUMENT TYPE: Journal

LANGUAGE: English


AB Measurements are reported on the kinetics of hydrolysis of dinitro-2-pyridylalanylglycine in solutions of hydrochloric acid up to 10M; acid-catalyzed hydrolysis of the amide bond occurs when the acid concentration is 2M or higher and the rate of hydrolysis is accurately proportional to H^+ ; classification of the reaction according to Bunnett gives an α value of zero and this indicates that water does not participate in transformation of SH^+ to a transition state. Furthermore the kinetic results for other dinitro-2-pyridyl dipeptides show that the catalytic effect is a general phenomenon. On the other hand, the hydrolysis of dinitro-2-pyridylalanylglycine is about 102-103 times as fast as the hydrolyses of dinitrophenylalanylglycine and of dinitro-4-pyridylalanylglycine. These results suggest that the acid-catalyzed hydrolysis of dinitro-2-pyridyl dipeptides is an electrophilic-nucleophilic catalyzed reaction involving a cyclic acylpyridinium intermediate.

IT 2900-30-3, Glycine, N-[1-(3,5-dinitro-2-pyridyl)prolyl]- (hydrolysis of)

RN 2900-30-3 HCAPLUS

CN Glycine, N-[1-(3,5-dinitro-2-pyridyl)-L-prolyl]- (7CI, 8CI) (CA INDEX NAME)

Absolute stereochemistry.



CC 44 (Amino Acids, Peptides, and Proteins)

IT 2900-13-2, Alanine, N-[N-(3,5-dinitro-2-pyridyl)alanyl]-
 2900-29-0, Alanine, N-[N-(3,5-dinitro-2-pyridyl)glycyl]-
 2900-30-3, Glycine, N-[1-(3,5-dinitro-2-pyridyl)prolyl]-
 2900-31-4, Serine, N-[N-(3,5-dinitro-2-pyridyl)glycyl]-
 2900-32-5, Leucine, N-[N-(3,5-dinitro-2-pyridyl)leucyl]-
 2900-33-6, Alanine, N-[N-(3,5-dinitro-2-pyridyl)alanyl]-3-phenyl-
 2900-34-7, Glycine, N-[N-(3,5-dinitro-2-pyridyl)alanyl]-
 2900-35-8, Glycine, N-[N-(2,4-dinitrophenyl)alanyl]- 2900-36-9,
 Glycine, N-[N-(3,5-dinitro-2-pyridyl)glycyl]-
 (hydrolysis of)

L26 ANSWER 10 OF 13 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1965:86288 HCAPLUS

DOCUMENT NUMBER: 62:86288

ORIGINAL REFERENCE NO.: 62:15402g-h

TITLE: Thin-layer chromatography of dinitropyridyl- and nitropyrimidylamino acids

AUTHOR(S): Di Bello, Carlo; Signor, Angelo

CORPORATE SOURCE: Univ. Padua, Italy

SOURCE: Journal of Chromatography (1965), 17(3), 506-12

CODEN: JOCRAM; ISSN: 0021-9673

DOCUMENT TYPE: Journal


LANGUAGE: English

AB One to 2 γ of sample was spotted on silica gel G coated plates. When the solvent front had travelled 10 cm., the plates

were removed and the solvent evaporated under a current of hot air. For 2-dimensional chromatog. the sample was applied onto the diagonal of the plate. For maximum reproducibility the plates must be dried between dimension runs at constant conditions. Spraying with 1% KMnO₄ followed by N HCl revealed nitropyrimidylamino acids as yellow spots on a pink background; they next appeared as black spots when sprayed with 8.1% HgNO₃ in 0.5N HNO₃, then 0.5N HNO₃, then, after drying, aqueous (NH₄)₂S. Of 6 solvent systems tested, either in single or 2-dimensional chromatog., CHCl₃-MeOH-AcOH (95:5:1) followed by ProH-33% NH₄OH (70:30) gave the best separation

IT 3264-09-3, Proline, 1-(3,5-dinitro-2-pyridyl)-, L-
(chromatog. of)
RN 3264-09-3 HCAPLUS
CN Proline, 1-(3,5-dinitro-2-pyridyl)-, L- (8CI) (CA INDEX NAME)

Absolute stereochemistry.



CC 2 (Analytical Chemistry)

IT 2980-33-8, 2-Pyridinol, 3,5-dinitro- 3073-24-3, Glutamic acid,
N-(3,5-dinitro-2-pyridyl)-, DL- 3073-25-4, Isoleucine,
N-(3,5-dinitro-2-pyridyl)-, DL- 3073-26-5, Leucine,
N-(3,5-dinitro-2-pyridyl)-, DL- 3073-27-6, Serine,
N-(3,5-dinitro-2-pyridyl)-, DL- 3073-28-7, Threonine,
N-(3,5-dinitro-2-pyridyl)-, DL- 3073-29-8, Valine,
N-(3,5-dinitro-2-pyridyl)-, DL- 3073-30-1, Pyridine,
2-amino-3,5-dinitro- 3073-32-3, Aspartic acid,
N-(5-nitro-2-pyrimidinyl)-, L- 3073-33-4, Glutamic acid,
N-(5-nitro-2-pyrimidinyl)-, DL- 3073-34-5, Alanine,
N-(5-nitro-2-pyrimidinyl)-, DL- 3073-68-5, Alanine,
N-(5-nitro-2-pyrimidinyl)-3-phenyl-, DL- 3073-69-6, Glycine,
N-(5-nitro-2-pyrimidinyl)-, L- 3073-70-9, Isoleucine,
N-(5-nitro-2-pyrimidinyl)-, DL- 3073-71-0, Leucine,
N-(5-nitro-2-pyrimidinyl)-, DL- 3073-72-1, Lysine,
N2,N6-bis(5-nitro-2-pyrimidinyl)-, L- 3073-73-2, Serine,
N-(5-nitro-2-pyrimidinyl)-, DL- 3073-74-3, Proline,
1-(5-nitro-2-pyrimidinyl)-, L- 3073-75-4, Threonine,
N-(5-nitro-2-pyrimidinyl)-, DL- 3073-76-5, Valine,
N-(5-nitro-2-pyrimidinyl)-, DL- 3073-77-6, Pyrimidine,
2-amino-5-nitro- 3264-06-0, Alanine, N-(3,5-dinitro-2-pyridyl)-,
DL- 3264-07-1, Alanine, N-(3,5-dinitro-2-pyridyl)-3-phenyl-, DL-
3264-08-2, Glycine, N-(3,5-dinitro-2-pyridyl)- 3264-09-3
, Proline, 1-(3,5-dinitro-2-pyridyl)-, L- 3264-10-6,
2-Pyrimidinol, 5-nitro- 3426-97-9, Lysine, N2,N6-bis(3,5-dinitro-
2-pyridyl)-, L- 3521-73-1, Aspartic acid, N-(3,5-dinitro-2-
pyridyl)-, L
(chromatog. of)

L26 ANSWER 11 OF 13 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1965:86287 HCAPLUS

DOCUMENT NUMBER: 62:86287

ORIGINAL REFERENCE NO.: 62:15402f-g

TITLE: Chromatographic separation of diketones. III.
Chromatography of derivatives of
2-phenyl-1,3-indandione

AUTHOR(S): Kreicberga, D.; Gudriniece, E.

SOURCE: Latvijas PSR Zinatnu Akademijas Vestis,
Kimijas Serija (1964), (4), 424-8

CODEN: LZAKAM; ISSN: 0002-3248

DOCUMENT TYPE:

Journal

LANGUAGE:

Russian


AB Derivs. (Cl, Br, I, NO₂, NH₂, Me₂N Et₂N, AcNH, MeO, SH, Me₃N, MeEt₂N) substituted in positions 2, 3, and 4 in the phenyl radical and in the 4 and 5 positions in the phthaloyl radical were tried. Best sepns. were achieved on powdered silica gel. Extensive tables, giving R_f values for 39 compds. in 12 different combinations of solvents and adsorbents are given.

IT 3264-09-3, Proline, 1-(3,5-dinitro-2-pyridyl)-, L- (chromatography of)

RN 3264-09-3 HCAPLUS

CN Proline, 1-(3,5-dinitro-2-pyridyl)-, L- (8CI) (CA INDEX NAME)

Absolute stereochemistry.



CC 2 (Analytical Chemistry)

IT 83-12-5, 1,3-Indandione, 2-phenyl 117-37-3, 1,3-Indandione, 2-(p-methoxyphenyl)- 606-23-5, 1,3-Indandione 969-63-1, 1,3-Indandione, 2-[p-(diethylamino)phenyl]- 1146-98-1, 1,3-Indandione, 2-(p-bromophenyl)- 1146-99-2, 1,3-Indandione, 2-(p-chlorophenyl)- 1147-00-8, 1,3-Indandione, 2-(p-iodophenyl)- 1153-90-8, 1,3-Indandione, 2-(p-nitrophenyl)- 1156-76-9, Acetanilide, 4'-(1,3-dioxo-2-indanyl)- 1225-30-5, 1,3-Indandione, 2-[p-(dimethylamino)phenyl]- 1470-34-4, 1,3-Indandione, 4-nitro-2-phenyl- 1470-35-5, 1,3-Indandione, 5-bromo-2-phenyl- 1470-36-6, 1,3-Indandione, 5-chloro-2-phenyl- 1470-37-7, 1,3-Indandione, 4-bromo-2-phenyl- 1470-38-8, 1,3-Indandione, 2-(3,4-dimethoxyphenyl)- 1470-39-9, 1,3-Indandione, 2-(o-methoxyphenyl)- 1470-40-2, 1,3-Indandione, 2-(o-iodophenyl)- 1470-41-3, 1,3-Indandione, 2-(o-bromophenyl)- 1470-42-4, 1,3-Indandione, 2-(o-chlorophenyl)- 1470-43-5, 1,3-Indandione, 2-(m-bromophenyl)- 1470-44-6, 1,3-Indandione, 2-(m-chlorophenyl)- 1470-52-6, 1,3-Indandione, 2-(p-(dimethylamino)phenyl)-4,5-dimethoxy- 1470-53-7, 1,3-Indandione, 4,5-dimethoxy-2-phenyl- 1470-54-8, 1,3-Indandione, 4-nitro-2-(p-nitrophenyl)- 1470-56-0, 1,3-Indandione, 4,5,6,7-tetrachloro-2-phenyl- 1640-36-4, 1,3-Indandione, 4-chloro-2-phenyl- 1640-37-5, 1,3-Indandione, 2-(m-iodophenyl)- 1641-12-9, 1,3-Indandione, 2-(3,4-dimethoxyphenyl)-4,5-dimethoxy- 1641-13-0, 1,3-Indandione, 4,5-dimethoxy-2-(p-methoxyphenyl)- 1641-14-1, 1,3-Indandione, 5-nitro-2-(p-nitrophenyl)- 1668-36-6, 1,3-Indandione, 2-(p-mercaptophenyl)- 1989-66-8, 1,3-Indandione, 4-iodo-2-phenyl- 2048-60-4, 1,3-Indandione, 5-iodo-2-phenyl- 2535-52-6, 1,3-Indandione, 5-nitro-2-phenyl- 2863-22-1, Acetanilide, 4'-(5-nitro-1,3-dioxo-2-indanyl)- 2878-47-9, Acetanilide, 4'-(4-nitro-1,3-dioxo-2-indanyl)- 2980-33-8, 2-Pyridinol, 3,5-dinitro- 3073-30-1, Pyridine, 2-amino-3,5-dinitro- 3073-68-5, Alanine, N-(5-nitro-2-pyrimidinyl)-3-phenyl-, DL- 3073-72-1, Lysine, N₂,N₆-bis(5-nitro-2-pyrimidinyl)-, L- 3073-74-3, Proline, 1-(5-nitro-2-pyrimidinyl)-, L- 3073-77-6, Pyrimidine, 2-amino-5-nitro- 3264-07-1, Alanine, N-(3,5-dinitro-2-pyridyl)-3-phenyl-, DL- 3264-09-3, Proline, 1-(3,5-dinitro-2-pyridyl)-, L- 3264-10-6, 2-Pyrimidinol, 5-nitro- 3426-97-9, Lysine, N₂,N₆-bis(3,5-dinitro-2-pyridyl)-, L- 3457-74-7,

1,3-Indandione, 4,5,6,7-tetrahydro-2-phenyl- 92497-72-8,
1,3-Indandione, 4-(p-aminophenyl)-
(chromatography of)

L26 ANSWER 12 OF 13 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1964:485898 HCAPLUS

DOCUMENT NUMBER: 61:85898

ORIGINAL REFERENCE NO.: 61:15018h,15019a-d

TITLE: A new method for the determination of
N-terminal amino acids in polypeptides and
proteins. III. Use of the reagent
2-chloro-3,5-dinitropyridine

AUTHOR(S): Signor, Angelo; Biondi, Laura; Terbojevich,
Maria; Pajetta, Paola

CORPORATE SOURCE: Univ. Padua

SOURCE: Gazzetta Chimica Italiana (1964), 94(6),
619-29

CODEN: GCITA9; ISSN: 0016-5603

DOCUMENT TYPE: Journal

LANGUAGE: Unavailable


AB cf. CA 61, 8611e. 2-Chloro-3,5-dinitropyridine (CA 59, 2811f) was found to react quant. with the N-terminal amino acids of proteins and peptides under relatively mild conditions to give the corresponding dinitropyridyl amino acids which were quant. recovered after acid hydrolysis. Further treatment of the dinitropyridyl amino acids with concentrated NH_4OH gave the free amino acids. Approx. 1 μmol of dialyzed and dried protein was dissolved or suspended in a solution consisting of an equal weight of NaHCO_3 and a 10-fold weight of H_2O . Two vols. of an alc. solution of 2-chloro-3,5-dinitropyridine were added and the solution was agitated at room temperature for 2 h., then acidified with concentrated HCl , and extracted with EtOAc . The recovered dinitropyridyl-protein was dissolved in 1 mL. 98-100% HCOOH and hydrolyzed in 6N HCl for 15-30 min. at 100° or for 10-15 h. at 60°. The latter time and temperature conditions were required for maximum recovery of the dinitropyridyl derivs. of serine, threonine, and proline which undergo partial destruction. The hydrolyzates were diluted with H_2O to pH 2 and extracted with 10-mL. portions of EtOAc . The several exts. were washed with 0.01N HCl and evaporated to dryness. The residue was dissolved in 5% NaHCO_3 and again extracted, after acidification, with EtOAc . The resulting dinitropyridyl amino acids were determined by ascending and descending quant. two-dimensional paper chromatog. (Levy, CA 49, 101h; Biserte and Osteux, CA 45 7622g). The spots were localized under UV light, and were each cut out and extracted with 4 mL. 1% NaHCO_3 at 60° for 30 min. After 10 min. at room temperature, the solns. were read at 340 $\text{m}\mu$. The dinitropyridyl proline was read at 360 $\text{m}\mu$. Data were obtained on the per cent amino acid liberated vs. hydrolysis time from the dinitropyridyl derivs. of 18 of the commonly occurring amino acids and glycylglycine, alanyl glycine, alanylphenylalanine, and leucylleucine. It is concluded that the principal advantages of the method are in the relatively mild conditions of forming the dinitropyridyl-proteins and the stability of the dinitropyridyl amino acids during hydrolysis.

IT 2900-30-3, Glycine, N-[1-(3,5-dinitro-2-pyridyl)-L-prolyl]-
3264-09-3, Proline, 1-(3,5-dinitro-2-pyridyl)-, L-
(in determination of amino acids in peptides)

RN 2900-30-3 HCAPLUS

CN Glycine, N-[1-(3,5-dinitro-2-pyridyl)-L-prolyl]- (7CI, 8CI) (CA
INDEX NAME)


Absolute stereochemistry.



RN 3264-09-3 HCAPLUS

CN Proline, 1-(3,5-dinitro-2-pyridyl)-, L- (8CI) (CA INDEX NAME)

Absolute stereochemistry.



CC 60 (Biochemical Methods)

IT 2578-45-2, Pyridine, 2-chloro-3,5-dinitro- 2900-30-3,
 Glycine, N-[1-(3,5-dinitro-2-pyridyl)-L-prolyl]- 2900-34-7,
 Glycine, N-[N-(3,5-dinitro-2-pyridyl)-DL-alanyl]- 2900-36-9,
 Glycine, N-[N-(3,5-dinitro-2-pyridyl)glycyl]- 3073-24-3,
 Glutamic acid, N-(3,5-dinitro-2-pyridyl)-, DL- 3073-25-4,
 Isoleucine, N-(3,5-dinitro-2-pyridyl)-, DL- 3073-26-5, Leucine,
 N-(3,5-dinitro-2-pyridyl)-, DL- 3073-27-6, Serine,
 N-(3,5-dinitro-2-pyridyl)-, DL- 3073-28-7, Threonine,
 N-(3,5-dinitro-2-pyridyl)-, DL- 3073-29-8, Valine,
 N-(3,5-dinitro-2-pyridyl)-, DL- 3264-06-0, Alanine,
 N-(3,5-dinitro-2-pyridyl)-, DL- 3264-07-1, Alanine,
 N-(3,5-dinitro-2-pyridyl)-3-phenyl-, DL- 3264-08-2, Glycine,
 N-(3,5-dinitro-2-pyridyl)- 3264-09-3, Proline,
 1-(3,5-dinitro-2-pyridyl)-, L- 3426-97-9, Lysine,
 N2,N6-bis(3,5-dinitro-2-pyridyl)-, L- 3521-73-1, Aspartic acid,
 N-(3,5-dinitro-2-pyridyl)-, L- 19339-97-0, Glutamine,
 N2-(3,5-dinitro-2-pyridyl)-, L- 91085-85-7, Asparagine,
 N2-(3,5-dinitro-2-pyridyl)-, L- 92546-66-2, Arginine,
 N2-(3,5-dinitro-2-pyridyl)-, L- 92872-58-7, Tryptophan,
 N-(3,5-dinitro-2-pyridyl)-, L- 94063-05-5, Leucine,
 N-[N-(3,5-dinitro-2-pyridyl)-DL-leucyl]-, DL- 94729-28-9,
 Alanine, N-[N-(3,5-dinitro-2-pyridyl)-DL-alanyl]-3-phenyl-, DL-
 (in determination of amino acids in peptides)

L26 ANSWER 13 OF 13 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1963:462817 HCAPLUS

DOCUMENT NUMBER: 59:62817

ORIGINAL REFERENCE NO.: 59:11650h,11651a-c

TITLE: Structure of proteins. VII. Preparation of nitropyridylamino acids

AUTHOR(S): Signor, Angelo; Scoffone, Ernesto; Biondi, Laura

CORPORATE SOURCE: Univ. Padua, Italy

SOURCE: Gazzetta Chimica Italiana (1963), 93(1-2), 73-80

CODEN: GCITA9; ISSN: 0016-5603

DOCUMENT TYPE: Journal

LANGUAGE: Unavailable


OTHER SOURCE(S): CASREACT 59:62817

AB cf. CA 59, 2811f. Adding 1.2 + 10⁻³ mole freshly prepared 2-

or 4-chloro-3,5-dinitropyridine in 40 cc. Me₂CO to 10⁻³ mole amino acid in 30 cc. 5% aqueous NaHCO₃, allowing the mixture to react 2-3 h. at ambient temperature with occasional stirring, evaporating solvent, diluting with H₂O to the original volume, extracting with CHCl₃, acidifying with 2N HCl to pH 2-3, filtering, and crystallizing from H₂O or H₂O-EtOH gave the 3,5-dinitro-4-(I) and 2-pyridylamino acids (II) in 80-98% yield, separated and identified by ascending paper chromatog. in 5:1:3:3 MePh-pyridine-Cl(CH₂)₂OH-0.8M NH₃ and determined at 380-90 mμ for the 4-pyridyl and at 340 ± 2 mμ (except proline derivative 360 mμ) for the 2-pyridyl derivative (amino acid and m.p. of I and II given): glycine, 192°, 165°; DL-alanine, 185°, 164°; DL-serine, 225° (decomposition), 153°; DL-valine, 183°, 167°; L-leucine, 166°, 104°; L-tyrosine, 160°, 107°; DL-phenylalanine, 205°, 148°; L-proline, 150°, 157°; L-glutamic acid, 215° (decomposition), -; D-lysine, 113°, -. The stability of the derivs. under conditions of acid protein hydrolysis was determined by heating 5 + 10⁻⁵ mole 16 h. at 110° in closed vials in 1 cc. 6N HCl, evaporating HCl, bringing to pH 2.2 with citric buffer, diluting to 50 cc., and determining a 2-cc. sample on an automatic analyzer. II showed much greater stability than I; e.g. 3,5-dinitro-2-pyridylvaline hydrolyzed only 2%. 2-Chloro-3,5-dinitropyridine is recommended as a reagent for terminal amino acid determination


IT 3264-09-3, Proline, 1-(3,5-dinitro-2-pyridyl)-, L-
(preparation of)
RN 3264-09-3 HCAPLUS
CN Proline, 1-(3,5-dinitro-2-pyridyl)-, L- (8CI) (CA INDEX NAME)

Absolute stereochemistry..



CC 44 (Amino Acids, Peptides, and Proteins)
IT 3073-26-5, Leucine, N-(3,5-dinitro-2-pyridyl)-, DL- 3073-27-6,
Serine, N-(3,5-dinitro-2-pyridyl)-, DL- 3073-29-8, Valine,
N-(3,5-dinitro-2-pyridyl)-, DL- 3264-06-0, Alanine,
N-(3,5-dinitro-2-pyridyl)-, DL- 3264-07-1, Alanine,
N-(3,5-dinitro-2-pyridyl)-3-phenyl-, DL- 3264-08-2, Glycine,
N-(3,5-dinitro-2-pyridyl)- 3264-09-3, Proline,
1-(3,5-dinitro-2-pyridyl)-, L- 82530-77-6, Glutamic acid,
N-(3,5-dinitro-2-pyridyl)-, L- 89677-25-8, Glycine,
N-(3,5-dinitro-4-pyridyl)- 89977-92-4, Alanine,
N-(3,5-dinitro-4-pyridyl)-, DL- 89977-97-9, Serine,
N-(3,5-dinitro-4-pyridyl)-, DL- 90871-34-4, Valine,
N-(3,5-dinitro-4-pyridyl)-, DL- 93350-89-1, Leucine,
N-(3,5-dinitro-4-pyridyl)-, DL-
(preparation of)


=> d que stat 131
L12 STR



VAR G2=C/N/O
 REP G3=(0-2) C
 NODE ATTRIBUTES:
 DEFAULT MLEVEL IS ATOM
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
 RING(S) ARE ISOLATED OR EMBEDDED
 NUMBER OF NODES IS 14

STEREO ATTRIBUTES: NONE
 L14 102 SEA FILE=REGISTRY SSS FUL L12
 L19 STR



VAR G1=NH2/NO2
 VAR G2=C/N/O
 REP G3=(0-2) C
 VAR G4=X/AK
 NODE ATTRIBUTES:
 DEFAULT MLEVEL IS ATOM
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
 RING(S) ARE ISOLATED OR EMBEDDED
 NUMBER OF NODES IS 15

STEREO ATTRIBUTES: NONE
 L21 53 SEA FILE=REGISTRY SUB=L14 SSS FUL L19
 L26 13 SEA FILE=HCAPLUS ABB=ON PLU=ON L21
 L28 28 SEA FILE=HCAPLUS ABB=ON PLU=ON L14
 L31 15 SEA FILE=HCAPLUS ABB=ON PLU=ON L28 NOT L26

=> d l31 1-15 ibib abs hitstr hitind

L31 ANSWER 1 OF 15 HCAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 2004:291093 HCAPLUS
 DOCUMENT NUMBER: 140:326608
 TITLE: New trinuclear heteroaromatic direct black dyes

INVENTOR(S): Kravtchenko, Sylvain; Lagrange, Alain; David, Herve; Greaves, Andrew; Bonaventure, Nicole; Vidal, Laurent
 PATENT ASSIGNEE(S): L'Oreal, Fr.
 SOURCE: Fr. Demande, 49 pp.
 CODEN: FRXXBL
 DOCUMENT TYPE: Patent
 LANGUAGE: French
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 2845387	A1	20040409	FR 2002-12385	2002 1004
FR 2845387	B1	20050121		
WO 2004031173	A1	20040415	WO 2003-FR2927	2003 1006
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2003283499	A1	20040423	AU 2003-283499	2003 1006
US 2005060815	A1	20050324	US 2003-678635	2003 1006
EP 1551825	A1	20050713	EP 2003-775473	2003 1006
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
PRIORITY APPLN. INFO.:		FR 2002-12385	A	2002 1004
		US 2002-431749P	P	2002 1209
		WO 2003-FR2927	W	2003 1006

OTHER SOURCE(S): MARPAT 140:326608

AB New trinuclear heteroarom. direct dyes, the hair dye compns. containing these dyes as well as the process of dyeing of keratinous fibers are disclosed. The trinuclear heteroarom. direct black dyes comprise a pyridin core. The new dyes make it possible to obtain black nuances which have a good tenacity and stability in the hair dye compns. Thus, 3-amino-7-diethylaminopyrazolopyrimidine was reacted with 2-pyrrolidino-3-amino-6-methoxypyridine to obtain a trinuclear


heteroarom. direct dye. Formulation of a dye containing 0.5% of above dye is disclosed.

IT 677314-81-7 677314-82-8 677314-83-9
677314-84-0 677314-85-1 677314-86-2
677314-87-3

RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(new trinuclear heteroarom. direct black dyes)


RN 677314-81-7 HCAPLUS

CN Pyrazolo[1,5-a]pyridin-3-amine, N-[2,5-dihydro-5-imino-2-[(2-methylpyrazolo[1,5-a]pyridin-3-yl)imino]-6-(1-pyrrolidinyl)-3-pyridinyl]-2-methyl- (9CI) (CA INDEX NAME)




RN 677314-82-8 HCAPLUS

CN Pyrimidinetetramine, N5-[2,5-dihydro-5-imino-6-(4-methyl-1-piperazinyl)-2-[(2,4,6-triamino-5-pyrimidinyl)imino]-3-pyridinyl]- (9CI) (CA INDEX NAME)




RN 677314-83-9 HCAPLUS

CN Pyrazolo[1,5-a]pyrimidine-3,7-diamine, N3-[3-[(7-amino-5-methylpyrazolo[1,5-a]pyridin-3-yl)amino]-5-imino-6-(1-pyrrolidinyl)-2(5H)-pyridinylidene]-5-methyl- (9CI) (CA INDEX NAME)




RN 677314-84-0 HCAPLUS

CN 3-Pyrrolidinaminium, 1-[3-amino-5,6-bis[[7-(dimethylamino)pyrazolo[1,5-a]pyrimidin-3-yl]imino]-5,6-dihydro-2-pyridinyl]-N,N,N-trimethyl-, chloride (9CI) (CA INDEX NAME)




RN 677314-85-1 HCAPLUS

CN Pyrazolo[1,5-a]pyrimidine-3,7-diamine, N3,N3'-[5-amino-6-[3-(dimethylamino)-1-pyrrolidinyl]-2,3-pyridinediylidene]bis[N7,N7-dimethyl- (9CI) (CA INDEX NAME)




RN 677314-86-2 HCAPLUS

CN Pyrazolo[1,5-a]pyridin-3-amine, N-[6-[3-(dimethylamino)-1-pyrrolidinyl]-2,5-dihydro-5-imino-2-(pyrazolo[1,5-a]pyridin-3-ylimino)-3-pyridinyl]- (9CI) (CA INDEX NAME)



RN 677314-87-3 HCAPLUS

CN 3-Pyrrolidinaminium, 1-[3,6-dihydro-3-imino-5-(pyrazolo[1,5-a]pyridin-3-ylamino)-6-(pyrazolo[1,5-a]pyridin-3-ylimino)-2-pyridinyl]-N,N,N-trimethyl- (9CI) (CA INDEX NAME)




IT 677314-72-6P 677314-80-6P

RL: COS (Cosmetic use); SPN (Synthetic preparation); BIOL
(Biological study); PREP (Preparation); USES (Uses)
(new trinuclear heteroarom. direct black dyes)


RN 677314-72-6 HCAPLUS

CN Pyrazolo[1,5-a]pyrimidine-3,7-diamine, N3,N3'-(5-amino-6-(1-pyrrolidinyl)-2,3-pyridinediylidene)bis[N7,N7-dimethyl- (9CI) (CA INDEX NAME)



RN 677314-80-6 HCAPLUS

CN 1H-Pyrazole-4,5-diamine, N4,N4'-(5-amino-6-(1-pyrrolidinyl)-2,3-pyridinediylidene)bis[1-ethyl-3-methyl- (9CI) (CA INDEX NAME)





IC ICM C07D471-04
ICS C07D401-14; A61K007-13; C07D231-38; C07D213-02; C07D239-42;
C07D213-74; C07D209-40; C07D241-04; C07D213-90
CC 62-3 (Essential Oils and Cosmetics)
Section cross-reference(s): 28
IT 677314-81-7 677314-82-8 677314-83-9
677314-84-0 677314-85-1 677314-86-2
677314-87-3 677314-88-4 677314-89-5
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(new trinuclear heteroarom. direct black dyes)
IT 677314-72-6P 677314-74-8P 677314-75-9P 677314-76-0P
677314-77-1P 677314-78-2P 677314-79-3P 677314-80-6P
RL: COS (Cosmetic use); SPN (Synthetic preparation); BIOL
(Biological study); PREP (Preparation); USES (Uses)
(new trinuclear heteroarom. direct black dyes)

L31 ANSWER 2 OF 15 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2001:98457 HCAPLUS
DOCUMENT NUMBER: 134:147611
TITLE: Preparation of tetrahydrobenzo[d]azepines as
metabotropic glutamate receptor 1 antagonists
INVENTOR(S): Adam, Geo; Binggeli, Alfred; Maerki,
Hans-Peter; Mutel, Vincent; Wilhelm, Maurice;
Wostl, Wolfgang
PATENT ASSIGNEE(S): F. Hoffmann-La Roche A.-G., Switz.
SOURCE: Eur. Pat. Appl., 85 pp.
CODEN: EPXXDW
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1074549	A2	20010207	EP 2000-116091	2000 0727
EP 1074549	A3	20020731		
EP 1074549	B1	20031119		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
AT 254614	E	20031215	AT 2000-116091	2000 0727
ES 2209728	T3	20040701	ES 2000-116091	2000

CA 2314798	AA	20010206	CA 2000-2314798	0727
				2000
				0801
US 6218385	B1	20010417	US 2000-630702	
				2000
				0801
NZ 506096	A	20020828	NZ 2000-506096	
				2000
				0801
ZA 2000003927	A	20010206	ZA 2000-3927	
				2000
				0802
AU 2000048979	A5	20010208	AU 2000-48979	
				2000
				0802
AU 774485	B2	20040701		
HR 2000000520	A1	20010630	HR 2000-520	
				2000
				0802
SG 93251	A1	20021217	SG 2000-4344	
				2000
				0802
NO 2000003966	A	20010207	NO 2000-3966	
				2000
				0804
CN 1283623	A	20010214	CN 2000-122523	
				2000
				0804
TR 200002298	A2	20010321	TR 2000-200002298	
				2000
				0804
JP 2001089472	A2	20010403	JP 2000-236848	
				2000
				0804
JP 3260350	B2	20020225		
RU 2240317	C2	20041120	RU 2000-120522	
				2000
				0804
BR 2000003375	A	20010313	BR 2000-3375	
				2000
				0807
PRIORITY APPLN. INFO.:			EP 1999-115557	A
				1999
				0806
OTHER SOURCE(S):		MARPAT 134:147611		
GI				



AB The title compds. (I) [wherein R1 = H, alkyl, O, halo, OR, cycloalkoxy, (un)substituted cycloalkylalkoxy, cyanoalkoxy, (fluoro)alkoxy, aminoalkoxy, alkenyloxy, phenylalkoxy, heterocyclalkoxy, sulfonyloxyalkoxy, SR, carboxyalkylthio, NR2, hydroxyalkylamino, or heterocyclalkylamino; n = 1-6; R = independently H, alkyl, or alkenyl; R2 = NO2 or CN; R3 = H, alkyl, O, S, SR, alkylsulfonyl, cycloalkyl, CONR2, NR2, alkyl, OR, or (un)substituted piperazino, carbamoylalkyl, alkoxyalkyl, fluoroalkyl, trifluoroacetoxyalkyl, carboxyalkyl, phenylthioalkyl, heterocyclalkoxy, acylamino, alkylamino, phenoxyalkylamino, heterocyclalkylamino, fluoroalkoxy, etc.; R4 = H, alkyl, alkenyl, NO2, OR, NR2, or (un)substituted fluoroalkoxy, fluoroalkyl, phenylalkyl, alkoxyalkanol, aminoalkyl, carboxyalkyl, alkylsulfonyloxyalkyl, fluoroalkenyl, heterocyclalkyl, heterocyclalkylamino, alkoxy carbonylamino, alkoxy carbonylhydrazino, aminofluoroalkenylamino; or R4 and R1 or R3 and R4 form an addnl. ring; R5 and R6 = independently H, alkyl, alkoxy, NH2, HO2, SO2NH2, or halo; or R5 and R6 = OCH2O; R7 and R8 = independently H, alkyl, alkoxy, NH2, NO2, or halo; R9 and R10 = independently H or alkyl; R11 and R12 = independently H, alkyl, OH, alkoxy, alkoxy carbonyloxy, or alkanoyloxy; R13 and R14 = independently H, T, or alkyl; R15 and R16 = independently H, T, alkyl, OH, alkoxy, alkoxy carbonyloxy, or alkanoyloxy; or R15 and R16 = O; X = N or C; Y = N, NH, or CH] were prepared. For example, addition of Et 2-cyano-3,3-bis(methylthio)acrylate to 2,3,4,5-tetrahydro-1H-benzo[d]azepine•HCl using TEA and K2CO3 in EtOH gave 2-cyano-3-methylsulfanyl-3-(1,2,4,5-tetrahydrobenzo[d]azepin-3-yl)acrylic acid Et ester (64%). The benzazepinylacrylate ester was treated with NH2C(NH)NH2•HNO3 and 1,8-diazabicyclo[5.4.0]undec-7-ene in DMF to give II (R = H). Ethylation of II (R = H) with EtI in DMF in the presence of K2CO3 afforded the preferred metabotropic glutamate receptor 1 (mGluR1) antagonist II (R = Et), which gave an IC50 values of 0.009 μ M and 0.003 μ M, resp. in functional and binding assays for the characterization of mGluR1 antagonist properties. I are useful in the prevention or control of acute and/or chronic neurol. disorders and as radiolabeled mGluR1 receptor antagonists in binding assays (no data).


IT 324554-07-6P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of tetrahydrobenzo[d]azepine mGluR1 antagonists by addition of chloroheterocycles or halobenzenes to tetrahydrobenzo[d]azepines or by cycloaddn. of guanidines to 3-methylthio-3-(tetrahydrobenzo[d]azepin-3-yl)acrylates)

RN 324554-07-6 HCAPLUS

CN 1H-3-Benzazepine, 3-(3,5-dinitro-2-pyridinyl)-2,3,4,5-tetrahydro-(9CI) (CA INDEX NAME)



IC ICM C07D403-04

ICS A61K031-55; A61P043-00; C07D403-14; C07D401-04; C07D223-04; C07D405-14; C07D487-04; C07D401-14; C07F007-18; C07D471-04; G01N033-50

CC 28-16 (Heterocyclic Compounds (More Than One Hetero Atom))

Section cross-reference(s): 1

IT	324552-61-6P	324552-63-8P	324552-71-8P	324552-73-0P
	324552-75-2P	324552-77-4P	324552-79-6P	324552-81-0P
	324552-83-2P	324552-85-4P	324552-87-6P	324552-89-8P
	324552-95-6P	324552-99-0P	324553-07-3P	324553-11-9P
	324553-15-3P	324553-19-7P	324553-23-3P	324553-27-7P
	324553-35-7P	324553-37-9P	324553-39-1P	324553-41-5P
	324553-43-7P	324553-45-9P	324553-47-1P	324553-49-3P
	324553-51-7P	324553-52-8P	324553-58-4P	324553-60-8P
	324553-62-0P	324553-63-1P	324553-67-5P	324553-69-7P
	324553-70-0P	324553-71-1P	324553-73-3P	324553-75-5P
	324553-77-7P	324553-78-8P	324553-80-2P	324553-84-6P
	324553-86-8P	324553-88-0P	324553-90-4P	324553-92-6P
	324553-94-8P	324553-96-0P	324553-98-2P	324554-00-9P
	324554-01-0P	324554-03-2P	324554-05-4P	324554-07-6P
	324554-09-8P	324554-11-2P	324554-13-4P	324554-15-6P
	324554-17-8P	324554-19-0P	324554-21-4P	324554-23-6P
	324554-30-5P	324554-32-7P	324554-38-3P	324554-40-7P
	324554-42-9P	324554-44-1P	324554-46-3P	324554-48-5P
	324554-52-1P	324554-54-3P	324554-56-5P	324554-58-7P
	324554-60-1P	324554-62-3P	324554-66-7P	324554-68-9P
	324554-70-3P	324554-72-5P	324554-74-7P	324554-76-9P
	324554-78-1P	324554-82-7P	324554-86-1P	324554-88-3P
	324554-90-7P	324554-92-9P	324554-94-1P	324554-96-3P
	324555-04-6P	324555-06-8P	324555-14-8P	324555-16-0P
	324555-18-2P	324555-20-6P	324555-24-0P	324555-26-2P
	324555-28-4P	324555-30-8P	324555-32-0P	324555-34-2P
	324555-38-6P	324555-40-0P	324555-42-2P	324555-44-4P
	324555-46-6P	324555-48-8P	324555-50-2P	324555-52-4P
	324555-56-8P	324555-58-0P	324555-64-8P	324555-66-0P
	324555-70-6P	324555-71-7P	324555-73-9P	324555-75-1P
	324555-77-3P	324555-83-1P	324555-87-5P	324555-89-7P
	324555-91-1P	324555-93-3P	324555-95-5P	324555-99-9P
	324556-01-6P	324556-03-8P	324556-07-2P	324556-09-4P
	324556-11-8P	324556-13-0P	324556-17-4P	324556-21-0P
	324556-23-2P	324556-25-4P	324556-29-8P	324556-31-2P
	324556-33-4P	324556-35-6P	324556-37-8P	324556-39-0P
	324556-41-4P	324556-43-6P	324556-45-8P	324556-47-0P
	324556-49-2P	324556-51-6P	324556-53-8P	324556-55-0P
	324556-57-2P	324556-59-4P	324556-61-8P	324556-63-0P